
A LIMIT THEOREM ON CHARACTERISTIC FUNCTIONS VIA AN EXTREMAL PRINCIPLE(U) TEXAS UNIV AT AUSTIN CENTER FOR CYBERNETIC STUDIES A BEN-TAL DEC 83 CCS-RR-477 F/G 12/1 AD-A141 390 1/1 -NL UNCLASSIFIED £



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

Research Report CCS 477

A LIMIT THEOREM ON CHARACTERISTIC FUNCTIONS

VIA AN EXTREMAL PRINCIPLE

bу

A. Ben-Tal\*

# CENTER FOR CYBERNETIC STUDIES

The University of Texas Austin, Texas 78712



This document has been approved for public release and sale: its distribution is unlimited.



84 05 21 148

# Research Report CCS 477 A LIMIT THEOREM ON CHARACTERISTIC FUNCTIONS VIA AN EXTREMAL PRINCIPLE

by

A. Ben-Tal\*

December 1983



\*Faculty of Industrial Engineering & Management, Technion-Israel Institute of Technology, Haifa, Israel

This paper is supported partly by ONR Contract N00014-81-C-0236 with the Center for Cybernetic Studies, The University of Texas at Austin and NSF Grant No. ECS-8214081 and Technion VPR Fund. Reproduction in whole or in part is permitted for any purpose of the United States Government.

### CENTER FOR CYBERNETIC STUDIES

A. Charnes, Director
Graduate School of Business 4.138
The University of Texas at Austin
Austin, Texas 78712
(512) 471-1821

This document has been approved for public release and sale; its distribution is unlimited.

### Abstract

We prove a classical limit theorem on characteristic functions by using duality between a pair of optimization problems, one of which is an infinite dimensional minimization involving the relative entropy functional.

KEY WORDS: Characteristic Functions, Optimization in infinite dimensional spaces, Duality, Relative Entropy.



### Introduction

Modern optimization theory has been employed successfully in many diverse fields such as Economics, Physics, Statistics, Biology and Engineering. This paper is a small step toward demonstrating the use of optimization theory as proof mechanism in Probability.

The result from Probability Theory in question here is a limit theorem on characteristic functions. Let X be a random variable with distribution  $\mathbf{F}_{\mathbf{X}}$ , support  $[\mathbf{x}_{\mathbf{L}},\mathbf{x}_{\mathbf{R}}]$  and characteristic function  $\psi(t) = \mathbf{E}\mathbf{e}^{itX}$ 

Then

$$x_R = \lim_{y \to \infty} \frac{1}{y} \log \psi(-iy)$$

$$\mathbf{x_L} = -\lim_{\mathbf{y} \to \infty} \frac{1}{\mathbf{y}} \log \psi(\mathbf{i}\mathbf{y}) .$$

The result is given in Lucas' classical book "Characteristic Functions"

[3]; first a weaker result, concerning only analytic characteristic functions, is proved in Chapter 7. The full statement is given in Chapter 11, as part of Theorem 11.1.2. It is derived from the result in Ch. 7 via a chain of lemmas on boundary characteristic functions.

Here we prove the above limit theorem by using duality relations between two extremum problems. One of these problems is an infinite-dimensional convex program involving the minimization of the relative entropy functional, which is of dundamental importance in Statistical Information Theory, Thermodynamics and Communication Theory.

The plan of the paper is as follows: Section 1 gives a formal statement of the limit theorem (Theorem A). Section 2 gives the duality theorem (Theorem B), which is in fact an adaptation of a result in the authors' paper [1]. In Section 3 we prove Theorem A via Theorem B.

### 1. A limit theorem on characteristic functions

Let X be a random variable, and  $F_{X}(x)$  its distribution function. Let  $\psi(t)$  denote the characteristic function of  $F_{X}$ , i.e.

$$\psi(t) = \int_{-\infty}^{\infty} e^{itx} dF_{X}(x) .$$

The left extremity of  $F_x$  is the number  $x_L$  with the property

$$\forall \epsilon > 0 : F_X(x_L - \epsilon) = 0 , F_X(x_L + \epsilon) > 0$$
.

and the right extremity of F is the number  $X_R$  with the property

$$\forall \epsilon > 0 : F_{\chi}(x_R - \epsilon) < 1 \qquad F_{\chi}(x_R) = 1$$

The interval [x,x,] is the support of Fx. Clearly

$$\psi(t) = \int_{x_L}^{x_R} e^{itx} dF_X(x) .$$

 $F_X$  is bounded to the left if  $x_L > -\infty$  and bounded to the right if  $x_R < \infty$  .

Theorem A. If F<sub>X</sub> be bounded to the right, then its right extremity is given by

(1) 
$$x_{R} = \lim_{y \to \infty} \frac{1}{y} \log \psi(-iy)$$

If F is bounded from the left, then its left extremity is given by

(2) 
$$x_{L} = -\lim_{y \to \infty} \frac{1}{y} \log \psi(iy).$$

# 2. A duality theorem on relative entropies

Let D be the class of generalized densities  $f = \frac{dF}{dt}$  (Radon-Nikodym derivatives) of distribution functions F, on a given probability space, with support  $[x_L, x_R]$ . In particular  $F_X \in D$  and  $f_X = \frac{dF_X}{dt}$ 

is the corresponding density. The relative entropy (divergence, Kullback-Leibler distance) between  $F \in D$  and  $F_{\chi}$  is given by the quantity

$$I(f;f_{x}) = \int_{x_{t}}^{x_{R}} f(t) \log \left[\frac{f(t)}{f_{x}(t)}\right] dt.$$

It is well known that  $I(\cdot;f_X)$  is a nonnegative convex functional and is equal to zero if and only if  $f = f_X$  (a.e. with respect to dt) see [2].

A special case of a problem studied in [1,Ch.3] is the infinite-dimensional convex program

(E) 
$$\inf\{I(f,f_x): \int_{X_L}^{X_R} g(t)f(t)dt \ge a\}$$

where g(t) is a given summable function. It was shown in [1] that a dual problem is given by

(H) 
$$\sup_{\mathbf{y} \geq 0} \{ \mathbf{a} \mathbf{y} - \log \int_{\mathbf{x}}^{\mathbf{x}_{R}} \mathbf{e}^{\mathbf{y} \mathbf{g}(t)} \mathbf{f}_{\mathbf{x}}(t) dt \} .$$

Moreover, from Th. 1 in [1] the following duality relations hold between (E) and (H).

Theorem B. If (E) is feasible then inf(E) is attained, sup(H) is finite and

$$min(E) = sup(H)$$
.

## 3. Proof of Theorem A via Theorem B

First note that the trivial inequality

$$\forall y > 0 : \mathbf{E}\mathbf{e}^{YX} \leq \mathbf{e}^{Yx}R$$

implies

$$\lim_{y\to\infty}\frac{1}{y}\log E e^{yX} \leq x_R$$

i.e.

(3) 
$$\lim_{y\to\infty}\frac{1}{y}\log\psi(-iy)\leq x_{R}.$$

Consider now the problem

for some fixed  $\varepsilon > 0$  . This is a special case of problem (E) with g(t) = t ,  $a = x_R - \varepsilon$  . The dual is

$$\sup_{\mathbf{Y}} \{ \mathbf{y}(\mathbf{x}_{R} - \mathbf{\epsilon}) - \log \int_{\mathbf{X}}^{\mathbf{X}_{R}} \mathbf{e}^{\mathbf{Y}t} \mathbf{f}_{\mathbf{X}}(t) dt \}$$

$$\mathbf{y} \ge 0 \qquad \mathbf{x}_{L}$$

i.e.

$$(D_{\varepsilon}) \qquad \sup_{y \ge 0} \{y(x_{R}^{-\varepsilon}) - \log \psi(-iy)\}$$

Problem (E ) is clearly feasible for every  $\varepsilon > 0$  , and we infer from Theorem B:

$$\Rightarrow \sup_{\varepsilon} (D_{\varepsilon}) \ge \lim_{y \to \infty} \{y(x_{R} - \varepsilon) - \log \psi(-iy)\} =$$

$$= \lim_{y \to \infty} y[x_{R} - \varepsilon - \frac{1}{y} \log \psi(-iy)]$$

Now, for the limit to be finite, it is necessary that:

(4) 
$$x_{R} - \varepsilon \leq \lim_{y \to \infty} \frac{1}{y} \log \psi(-iy) \quad \forall \varepsilon > 0.$$

Combining (3) and (4) we obtain equation (1).

To prove equation (2) we note that the inequality

(5) 
$$-\lim_{y\to\infty}\frac{1}{y}\log\psi(iy)\geq x_{L}$$

is trivial, while the inequality

(6) 
$$-\lim_{Y\to\infty}\frac{1}{Y}\log\psi(iy)\leq x_L+\varepsilon \quad \forall \ \varepsilon>0$$

follows by applying Theorem B (in the above manner) to the dual pair:

$$\inf\{I(f,f_{x}): \int_{x_{L}}^{R} tf(t)dt \ge x_{L} - \epsilon\}$$

$$\sup\{-y(x_{L}+\epsilon) - \log \psi(iy)\}.$$

$$y \ge 0$$

Now, (5) and (6) indeed imply (2), and the proof of Theorem A is thereby completed.

### References

- [1] Ben-Tal, A., "The entropic penalty approach to stochastic programming", Math. of Operations Research (to appear).
- [2] Kullback, S., <u>Information Theory and Statistics</u>, Wiley, New York, 1959.
- [3] Lucas, E., Characteristic Functions (Second Edition), Griffin, London, 1970.

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
CCS 477	AD . A141390	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
A Limit Theorem on Characteristic Functions		
Via an Extremal Principle		6. PERFORMING ORG, REPORT NUMBER
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(#)
A. Ben-Tal		N00014-81-C-0236
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Center for Cybernetic Studies		
The University of Texas at Aust Austin, Texas 78712	in	
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Office of Naval Research (Code 434)		December 1983
Washington, D.C.		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS(It differen	it from Controlling Office)	15. SECURITY CLASS. (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
		SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
This document has been approved for public release and sale; its		
distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, If different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Characteristic functions, optimization in infinite dimensional spaces,		
duality, relative entropy		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
We prove a classical limit theorem on characteristic functions by using duality between a pair of optimization problems, one of which is		
an infinite dimensional minimization involving the relative entropy		
functional.		